

Nonlinear metrology with a quantum interface

M. Napolitano*, M. W. Mitchell

ICFO-Institut de Ciències Fòniques, 08860 Castelldefels (Barcelona), Spain

E-mail: *mario.napolitano@icfo.es

Abstract. We describe nonlinear quantum atom-light interfaces and nonlinear quantum metrology in the collective continuous variable formalism. We develop a nonlinear effective Hamiltonian in terms of spin and polarization collective variables and show that model Hamiltonians of interest for nonlinear quantum metrology can be produced in ^{87}Rb ensembles. With these Hamiltonians, metrologically relevant atomic properties, e.g. the collective spin, can be measured better than the “Heisenberg limit” $\propto 1/N$. In contrast to other proposed nonlinear metrology systems, the atom-light interface allows both linear and non-linear estimation of the same atomic quantities.

Submitted to: *New Journal of Physics*

PACS numbers: 42.50.Hz, 42.50.Dv, 42.65.An

1. Introduction

In quantum metrology, a quantum state is prepared, evolves under the action of a Hamiltonian containing a parameter x of interest, and is measured. The parameter is estimated from the measurement outcome and knowledge of the system and Hamiltonian. In most problems, the Hamiltonian is assumed to act in the same way on each of N systems (e.g. atoms), and precision scales as $\delta x \propto N^{-1/2}$ for product states, and down to $\delta x \propto N^{-1}$ for entangled states (“Heisenberg limit” scaling) [1]. A number of studies [2, 3, 4] have considered also nonlinear quantum metrology, in which the Hamiltonian describes a k -system coupling with strength x' . Remarkably, the scaling is $\delta x' \propto N^{-k+1/2}$ or $\delta x' \propto N^{-k}$ for independent or entangled states, respectively [2]. Because this improves upon the best possible scaling for the linear case, it has been called “Super-Heisenberg” (SH) scaling [4]. Proposed implementations include scattering in Bose condensates [4], Duffing nonlinearity in nano-mechanical resonators [5], two-pass effective nonlinearity with an atomic ensemble [6], and Kerr nonlinearities [7, 8, 9].

Here we describe nonlinear metrology applied to measurement of collective spin variables of atomic ensembles. Atomic ensembles with long coherence-time internal degrees of freedom, e.g. nuclear spin, are essential elements of many quantum information and quantum metrology protocols including quantum memory [10], quantum non-demolition measurement [11], spin squeezing [12, 13, 14], and magnetometry [15].

We follow the approach of collective continuous variables (CCV), in which both light and atoms are described by macroscopic quantum variables. In the case of spinor atoms interacting with polarized light, the N_A atoms are described by the collective spin $\mathbf{F} \equiv \sum_i \mathbf{f}^{(i)}$ where $\mathbf{f}^{(i)}$ is the spin of the i -th atom. $\mathbf{F}$ obeys the commutation relations $[F_x, F_y] = i\hbar F_z$ and cyclic permutations, and can itself be considered a macroscopic spin variable. The light is described by its electric field $\mathbf{E} = \mathcal{E} + \mathcal{E}^*$, where $\mathcal{E}$ is the positive-frequency part. The Stokes vector $\mathbf{S}$ with components $S_i = (\mathcal{E}_+^*, \mathcal{E}_-^*) \sigma_i (\mathcal{E}_+, \mathcal{E}_-)^T$ where the subscript indicates plus/minus circular polarization, σ_i are the Pauli matrices and σ_0 is the identity. As described by several authors [16, 17, 18], the electric dipole interaction $h_{\text{int}} = -\mathbf{E} \cdot \mathbf{d}$, taken in second order perturbation theory, gives rise to an effective (single-atom) Hamiltonian of the form

$$h_{\text{eff}} = \sum_k \frac{\mathcal{E}^* \cdot \mathbf{d}_\downarrow |\phi_k\rangle \langle \phi_k| \mathbf{d}_\uparrow \cdot \mathcal{E}}{\hbar \delta_k} \quad (1)$$

$$= \mathcal{E}^* \cdot \overset{\leftrightarrow}{\alpha} \cdot \mathcal{E} \quad (2)$$

where $d_\uparrow, d_\downarrow$ are the parts of the dipole operator causing upward/downward transitions, δ_k is the detuning from resonance of the k th state, and $\overset{\leftrightarrow}{\alpha}$ is the tensor polarizability operator. The effective Hamiltonian for the ensemble $H_{\text{eff}} = \sum_i h_{\text{eff}}^{(i)}$ can then be decomposed into irreducible tensor components as

$$H_{\text{eff}}^{(2)} = \alpha_1 S_z J_z + \alpha_2 (S_x J_x + S_y J_y) , \quad (3)$$

plus terms in S_0 which do not interact with the optical polarization. Here $\alpha_{1,2}$ describe

the vectorial, and tensorial components of the interaction, respectively, and the atomic collective variable is $\mathbf{J} \equiv \sum_i \mathbf{j}^{(i)}$ with $j_x \equiv (f_x^2 - f_y^2)/2$, $j_y \equiv (f_x f_y + f_y f_x)/2$, $j_z \equiv f_z/2$ and $j_0 \equiv f_z^2/2$. The ratio of the α_i can be tuned by adjusting the optical frequency ω , giving a variety of Hamiltonians interesting for quantum information tasks [18].

To apply this formalism to nonlinear metrology, we generalize the CCV method to the nonlinear optics regime, i.e., we include higher-order processes in the effective Hamiltonian. For this purpose, naïve application of higher-order perturbation theory fails due to the appearance of vanishing resonance denominators, and degenerate perturbation theory [19] is required. We present the method by way of an example, the D_2 line of ^{87}Rb , one of the most used transitions for atom-light interactions.

2. Derivation of the Effective Hamiltonian

We consider the $5^2S_{1/2} \rightarrow 5^2P_{3/2}$ transition (the D_2 transition at 780 nm). The ground states are $|F, m_F\rangle$ with $(F, m_F) = (1, -1), (1, 0), (1, 1), (2, -2), \dots, (2, 2)$. The excited states are $|F', m_{F'}\rangle$ with $(F', m_{F'}) = (0, 0), (1, -1), (1, 0), \dots, (3, 3)$. We use these states as a basis, with the ground states preceding the excited states. We calculate the single-atom Hamiltonian h_{eff} , and note that the ensemble Hamiltonian $H_{\text{eff}} = \sum_i h_{\text{eff}}^{(i)}$ is found simply by replacing single-atom operators such as $\mathbf{j}$ with collective operators $\mathbf{J} \equiv \sum_i \mathbf{j}^{(i)}$. The unperturbed Hamiltonian is $h_0 = \hbar \sum_l \omega_l |l\rangle \langle l|$, or in matrix notation

$$h_0 = \hbar(\omega_{F=1} I_3 \oplus \omega_{F=2} I_5 \oplus \omega_{F'=0} I_1 \oplus \omega_{F'=1} I_3 \oplus \omega_{F'=2} I_5 \oplus \omega_{F'=3} I_7). \quad (4)$$

where $\oplus$ indicates a direct sum, and I_d is the identity matrix of dimension d . We choose the origin of energy such that $\omega_{F=1} = 0$, and define $\Delta \equiv \omega_{F=2}$. We work in a frame rotating with the laser frequency $\omega = \omega_{F'=0} + \delta$. In this frame, the Hamiltonian is

$$h_0 = \hbar(0 I_3 \oplus \Delta I_5 \oplus \delta_0 I_1 \oplus \delta_1 I_3 \oplus \delta_2 I_5 \oplus \delta_3 I_7) \quad (5)$$

where $\delta_{F'} \equiv \omega_{F'} - \omega$.

In the rotating wave approximation, the single-atom perturbation $v = h_{\text{int}} = -\mathbf{E} \cdot \mathbf{d}$ is approximated as $v \approx \mathcal{E} \cdot \mathbf{d}_{\uparrow} + \mathcal{E}^* \cdot \mathbf{d}_{\downarrow}$. If $\mathcal{E}_{\pm}$ are the amplitudes for the sigma-plus/minus components, respectively, of $\mathcal{E}$, then

$$\langle F', m_{F'} | V | F, m_F \rangle = \mathcal{E}_q \langle F', m_{F'} | er_q | F, m_F \rangle \quad (6)$$

with $q = m_{F'} - m_F$. Note that $q = 0$ transitions (π -transitions) are not considered because the z -propagating beam cannot contain this polarization. The dipole matrix elements are related to the “matrix element” $\langle J | |er_q| | J' \rangle \equiv D_{JJ'} \approx 3.58410^{-29} \text{C} \cdot \text{m}$ by angular-momentum addition rules. We follow the conventions given in Steck [20]. In this way, we arrive to the perturbation Hamiltonian

$$V = \begin{pmatrix} 0_8 & V_{\uparrow}^{\dagger} \\ V_{\uparrow} & 0_{16} \end{pmatrix} \quad (7)$$

where $0_d = 0I_d$ and $V_{\uparrow} \equiv \sqrt{5}D_{JJ'} \times$

$$\begin{pmatrix} \frac{\varepsilon_+}{\sqrt{30}} & 0 & \frac{\varepsilon_-}{\sqrt{30}} & 0 & 0 & 0 & 0 & 0 \\ 0 & \frac{\varepsilon_-}{2\sqrt{6}} & 0 & \frac{\varepsilon_+}{10} & 0 & \frac{\varepsilon_-}{10\sqrt{6}} & 0 & 0 \\ \frac{\varepsilon_+}{2\sqrt{6}} & 0 & \frac{\varepsilon_-}{2\sqrt{6}} & 0 & \frac{\varepsilon_+}{10\sqrt{2}} & 0 & \frac{\varepsilon_-}{10\sqrt{2}} & 0 \\ 0 & \frac{\varepsilon_+}{2\sqrt{6}} & 0 & 0 & 0 & \frac{\varepsilon_-}{10\sqrt{6}} & 0 & \frac{\varepsilon_-}{10} \\ \frac{\varepsilon_-}{2\sqrt{5}} & 0 & 0 & 0 & \frac{\varepsilon_-}{2\sqrt{15}} & 0 & 0 & 0 \\ 0 & \frac{\varepsilon_-}{2\sqrt{10}} & 0 & \frac{\varepsilon_+}{2\sqrt{15}} & 0 & \frac{\varepsilon_-}{2\sqrt{10}} & 0 & 0 \\ \frac{\varepsilon_+}{2\sqrt{30}} & 0 & \frac{\varepsilon_-}{2\sqrt{30}} & 0 & \frac{\varepsilon_+}{2\sqrt{10}} & 0 & \frac{\varepsilon_-}{2\sqrt{10}} & 0 \\ 0 & \frac{\varepsilon_+}{2\sqrt{10}} & 0 & 0 & 0 & \frac{\varepsilon_+}{2\sqrt{10}} & 0 & \frac{\varepsilon_-}{2\sqrt{15}} \\ 0 & 0 & \frac{\varepsilon_+}{2\sqrt{5}} & 0 & 0 & 0 & \frac{\varepsilon_+}{2\sqrt{15}} & 0 \\ 0 & 0 & 0 & \frac{\varepsilon_-}{\sqrt{10}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{\varepsilon_-}{\sqrt{15}} & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{\varepsilon_+}{5\sqrt{6}} & 0 & \frac{\varepsilon_-}{5} & 0 & 0 \\ 0 & 0 & 0 & 0 & \frac{\varepsilon_+}{5\sqrt{2}} & 0 & \frac{\varepsilon_-}{5\sqrt{2}} & 0 \\ 0 & 0 & 0 & 0 & 0 & \frac{\varepsilon_+}{5} & 0 & \frac{\varepsilon_-}{5\sqrt{6}} \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\varepsilon_+}{\sqrt{15}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \frac{\varepsilon_+}{\sqrt{10}} \end{pmatrix} \quad (8)$$

To obtain the effective Hamiltonian, we follow Klein [19]. The notation of that work is somewhat obscure, so for ease of understanding we repeat the main results. From Equation (A7) of that work, we have the t -order contribution to the effective Hamiltonian

$$h_{\text{eff}}^{(t)} = \sum_{\{k\}} A_{\{k\}} O_{\{k\}} \quad (9)$$

Where $k_1, \dots, k_{t-1}$ are non-negative integers, the A are real coefficients, the O , denoted “ $(k_1, k_2, \dots, k_t)$ ” by Klein, are operators, and the sum is taken over all $\{k\}$ satisfying $\sum_{l=1}^{t-1} k_l = t-1$. The A are given in Table I of that work and the O are given in Equation (A1) as

$$O_{\{k_1, \dots, k_{t-1}\}} \equiv P_0 V R^{(k_1)} V R^{(k_2)} \dots V R^{(k_{t-1})} V P_0 \quad (10)$$

with P_0 being the projector onto the degenerate subspace and by Equation (II.A.5)

$$R^{(k)} \equiv \begin{cases} P_0 & k = 0 \\ \left(\frac{1-P_0}{E_0-H^{(0)}} \right)^k & k > 0 \end{cases} \quad (11)$$

where E_0 is the energy of the degenerate subspace. In our case we have chosen $E_0 = 0$.

We can then directly calculate the second- and fourth-order contributions. We are only concerned with h_{eff} as it acts on the $F = 1$ subspace, that is, with a 3×3 matrix, and it is convenient to express it in terms of the pseudo-spin components j_0, j_x, j_y, j_z and the Stokes components S_0, S_x, S_y, S_z defined above. Summing the second-order




Figure 1. (color online). Spectra of the terms of the effective Hamiltonian. First two curves from the top, left axis: continuous, $\alpha^{(1)}$; dashed, $\alpha^{(2)}$. Lower curves, right axis: continuous, $\beta^{(1)}$; dotted, $\beta_N^{(0)}$; dashed, $\beta^{(2)}$; dot-dashed, $\beta_J^{(0)}$. Detuning [MHz] is relative to the transition $F = 1 \rightarrow F' = 0$ of ^{87}Rb D_2 transition. Points A, B, indicate detunings at which $\alpha^{(1)}$ or $\beta^{(1)}$ vanish.

contributions we find $H_{\text{eff}}^{(2)}$ of equation 3 With $B \equiv -D_{JJ'}^2/48\delta_0\delta_1\delta_2\hbar$,

$$\alpha^{(1)} = B(5\delta_0\delta_1 - 5\delta_0\delta_2 - 4\delta_1\delta_2) \quad (12)$$

$$\alpha^{(2)} = B(\delta_0\delta_1 - 5\delta_0\delta_2 + 4\delta_1\delta_2) \quad (13)$$

Similarly, the fourth-order contribution is, dropping terms in S_0^2 ,

$$H_{\text{eff}}^{(4)} = \beta_J^{(0)} S_Z^2 J_0 + \beta_N^{(0)} S_Z^2 N_A + \beta^{(1)} S_0 S_Z J_Z + \beta^{(2)} S_0 (S_X J_X + S_Y J_Y). \quad (14)$$

Note that the term in N_A arises because $h_{\text{eff}}^{(4)}$ contains a self-rotation term of the form $\beta_{m=0}^{(0)} S_Z^2 P_{m=0}$ where $P_{m=0}$ is a projector onto the state $|F = 1, m_F = 0\rangle$. We express this in terms of J_0 and N_A using $\sum_i P_{m=0}^{(i)} = \sum_i (I_3^{(i)} - j_0^{(i)}) = N_A - J_0$.

With $C \equiv D_{JJ'}^4/1152\delta_0^3\delta_1^3\delta_2^3\Delta\hbar^3$, the coefficients, shown graphically in Figure 1, are

$$\beta_J^{(0)} = C (12\delta_0^3\delta_1^2\delta_2^2 - 4\delta_0^3\delta_1\delta_2^3 + 12\delta_0^3\delta_1^3\Delta - 10\delta_0^3\delta_1^2\delta_2\Delta - 12\delta_0^2\delta_1^3\delta_2\Delta - 10\delta_0^3\delta_1\delta_2^2\Delta - 12\delta_0\delta_1^3\delta_2^2\Delta + 20\delta_0^2\delta_1\delta_2^3\Delta + 20\delta_0\delta_1^2\delta_2^3\Delta) \quad (15)$$

$$\beta_N^{(0)} = C (-12\delta_0^3\delta_1^3\delta_2 - 24\delta_0^3\delta_1^2\delta_2^2 + 4\delta_0^3\delta_1\delta_2^3) \quad (16)$$

$$\beta^{(1)} = C (-9\delta_0^3\delta_1^3\delta_2 + 6\delta_0^3\delta_1^2\delta_2^2 + 3\delta_0^3\delta_1\delta_2^3 + 35\delta_0^3\delta_1^3\Delta - 5\delta_0^3\delta_1^2\delta_2\Delta - 4\delta_0^2\delta_1^3\delta_2\Delta - 5\delta_0^3\delta_1\delta_2^2\Delta - 4\delta_0\delta_1^3\delta_2^2\Delta - 25\delta_0^3\delta_2^3\Delta - 20\delta_0^2\delta_1\delta_2^3\Delta - 20\delta_0\delta_1^2\delta_2^3\Delta - 16\delta_1^3\delta_2^3\Delta) \quad (17)$$

$$\begin{aligned}
\beta^{(2)} = C & (3\delta_0^3\delta_1^3\delta_2 - 6\delta_0^3\delta_1^2\delta_2^2 + 3\delta_0^3\delta_1\delta_2^3 + 7\delta_0^3\delta_1^3\Delta - 15\delta_0^3\delta_1^2\delta_2\Delta \\
& + 16\delta_0^2\delta_1^3\delta_2\Delta - 15\delta_0^2\delta_1\delta_2^2\Delta + 16\delta_0\delta_1^3\delta_2^2\Delta - 25\delta_0^3\delta_2^3\Delta \\
& + 16\delta_1^3\delta_2^3\Delta)
\end{aligned} \tag{18}$$

3. Application to nonlinear metrology

The β terms are nonlinear in $\mathbf{S}$, indicating a photon-photon interaction. We expect these terms to describe polarization effects of fast electronic nonlinearities including saturation and four-wave mixing. As in the linear case, the frequency dependence of the β terms provides considerable flexibility in designing a light-matter interaction. Applied to quantum metrology, these terms produce SH scaling, because they are nonlinear in the S collective variables, while the atomic variables J_i , N_A play the role of the parameter. The $\beta^{(0)}$ and $\beta^{(1)}$ terms are analogous to Hamiltonians considered by Boixo *et al.* [2]. The $\beta^{(1)}$ term $\propto S_0 S_z$, in particular, achieves SH scaling without input or generated entanglement [4]. The $\beta^{(2)}$ terms describes a nonlinear tensorial contribution, and does not appear to have been considered yet for nonlinear metrology.

4. Quantum Noise

To understand the quantum noise in this system, we define polarization operators $\hat{S}_i \equiv \frac{1}{2}(a_+^\dagger, a_-^\dagger)\sigma_i(a_+, a_-)^T$, where $a_\pm$ are annihilation operators for the $\pm$ circular polarizations of a mode defined by the pulse shape. These obey angular momentum commutation relations $[\hat{S}_i, \hat{S}_j] = i\varepsilon_{ijk}\hat{S}_k$ and are related to the Stokes parameters by $\hat{S}_i = S_i/2\gamma$, where $\gamma \equiv \hbar\omega Z_0/2TA$ is the single-photon intensity, T is the pulse duration, A is the beam area, and Z_0 is the impedance of free space. The total number of photons is $N_L = 2\hat{S}_0$. For a typical input, a coherent state, $\langle (\hat{S}_X, \hat{S}_Y, \hat{S}_Z) \rangle = (\hat{S}_0, 0, 0)$ and $\text{var}(\hat{S}_i) = \hat{S}_0/2$.

Evolution under this effective Hamiltonian produces, to first order in the interaction time τ ,

$$\hat{S}_Y^{(\text{out})} = \hat{S}_Y^{(\text{in})} + \frac{\tau}{\hbar}(\alpha^{(1)} + \beta^{(1)}\gamma\hat{S}_0)\gamma\hat{S}_X^{(\text{in})}J_Z^{(\text{in})} \tag{19}$$

plus terms containing $\hat{S}_Z^{(\text{in})}J_X^{(\text{in})}$ that are negligible for the given input coherent state of the light. This evolution physically corresponds to a paramagnetic Faraday rotation of the input linear polarization. In a metrological scheme one would measure this polarization rotation and from it estimate the atomic variable J_Z .

For small rotation, i.e. $\phi \equiv \hat{S}_Y^{(\text{out})}/\hat{S}_X^{(\text{in})} = \gamma\tau\hbar^{-1}(\alpha^{(1)} + \beta^{(1)}\gamma\hat{S}_0)J_Z^{(\text{in})} \ll 1$, we note that the input polarization noise dominates: $\text{var}(\hat{S}_Y^{(\text{out})}) = \text{var}(\hat{S}_Y^{(\text{in})}) + \phi^2\text{var}(\hat{S}_X) \approx \text{var}(\hat{S}_Y^{(\text{in})}) = \hat{S}_0/2$, and that the signal-to-noise ratio equals one when $\langle \hat{S}_Y^{(\text{out})} \rangle^2 = \text{var}(\hat{S}_Y^{(\text{in})})$, i.e., when

$$\frac{\tau^2\gamma^2}{\hbar^2}\hat{S}_0^2(\alpha^{(1)} + \beta^{(1)}\gamma\hat{S}_0)^2(J_Z^{(\text{in})})^2 = \frac{\hat{S}_0}{2}. \tag{20}$$

We can identify the value of $J_Z^{(\text{in})}$ that solves Eq. (20) as the sensitivity, or precision of the estimation, δJ_Z . We find

$$\delta J_Z = \hbar |\tau \gamma (\alpha^{(1)} N_L^{1/2} + \frac{\beta^{(1)} \gamma}{2} N_L^{3/2})|^{-1}. \quad (21)$$

Thus the sensitivity will have a transition from shot-noise to SH scaling with increasing N_L . As indicated in Figure 1, there are points in the spectrum where either $\alpha^{(1)}$ or $\beta^{(1)}$ vanish, allowing pure nonlinear or pure linear estimation of the same atomic variable.

In another scenario, an unpolarized input state $\langle (\hat{S}_X, \hat{S}_Y, \hat{S}_Z) \rangle = (0, 0, 0)$ gives rise to dynamics dominated by the $\beta^{(0)}$ terms $\propto S_Z^2$, sometimes called the “one-axis twisting Hamiltonian.” This describes a self-rotation of the optical polarization, and can be used to generate polarization squeezing and also to obtain sensitivity scaling as $N_L^{-3/2}$ in the estimation of $\beta_J^{(0)} J_0 + \beta_N^{(0)} N_A$, using an entanglement-generating strategy described in reference [4].

5. Conclusion

We have generalized the formalism of continuous collective variables to the nonlinear regime. The resulting nonlinear effective Hamiltonian includes several distinct nonlinear couplings with strengths widely tunable via the probe light frequency. This allows the production of model Hamiltonians proposed for nonlinear metrology, including both models that generate entanglement and those which achieve super-Heisenberg scaling without entanglement. Similar nonlinear probing techniques could improve optical probing of atomic clocks [21, 13] and atomic magnetometers [22, 15, 23]. Unlike previous proposals, the atomic ensemble system allows both linear and nonlinear estimation of the same atomic variables.

Acknowledgement

We thank M. Koschorreck, B. Dubost, N. Behbood, A. Cer  and R. Sewell for helpful discussion. F. Illuminati contributed essential ideas at an early stage. This work was funded by the Spanish Ministry of Science and Innovation under the FPU project and the Consolider-Ingenio 2010 Project “QOIT”

References

- [1] V. Giovannetti, S. Lloyd, and L. Maccone, *Phys. Rev. Lett.* **96**, 010401 (2006).
- [2] S. Boixo, S. T. Flammia, C. M. Caves, and J. Geremia, *Phys. Rev. Lett.* **98**, 090401 (2007).
- [3] S. Boixo et al., *Phys. Rev. A* **77**, 012317 (2008).
- [4] S. Boixo et al., *Phys. Rev. Lett.* **101**, 040403 (2008).
- [5] M. J. Woolley, G. J. Milburn, and C. M. Caves, *New J. Phys.* **10**, 125018 (2008).
- [6] B. A. Chase, B. Q. Baragiola, H. L. Partner, B. D. Black, and J. M. Geremia, *Phys. Rev. A* **79**, 062107 (2009).
- [7] A. Luis, *Phys. Lett. A* **329**, 8 (2004).
- [8] J. Beltr n and A. Luis, *Phys. Rev. A* **72**, 045801 (2005).

- [9] A. Luis, Phys. Rev. A **76**, 035801 (2007).
- [10] B. Julsgaard, J. Sherson, J. I. Cirac, J. Fiurášek and E. S. Polzik, Nature **432**, 482 (2007).
- [11] M. Koschorreck, M. Napolitano, B. Dubost, and M. W. Mitchell Phys. Rev. Lett. **102**, 033601 (2009).
- [12] P. J. Windpassinger, D. Oblak, U. B. Hoff, A. Louchet, J. Appel, N. Kjaergaard, E. S. Polzik J. Mod. Opt. **41**, 2315 (1994).
- [13] M. H. Schleier-Smith, I. D. Leroux, and V. Vuletić, Phys. Rev. Lett. **104**, 073604 (2010).
- [14] T. Takano, M. Fuyama, R. Namiki, and Y. Takahashi Phys. Rev. Lett. **104**, 093602 (2010).
- [15] D. Budker and M. Romalis, Nature Physics **3**, 227 (2007).
- [16] D. V. Kupriyanov, O. S. Mishina, and I. M. Sokolov, Phys. Rev. A **71**, 032348 (2005).
- [17] J. Geremia, J. Stockton, and H. Mabuchi, Phys. Rev. A **73**, 042112 (2006).
- [18] S. R. de Echaniz, M. Koschorreck, M. Napolitano, M. Kubasik, and M. W. Mitchell, Phys. Rev. A **77**, 032316 (2008).
- [19] D. J. Klein, J. Chem. Phys. **61**, 786 (1974).
- [20] D. A. Steck, Rubidium 87 D line data, <http://steck.us/alkalidata>, 2009.
- [21] J. Appel et al., PNAS **106**, 10960 (2009).
- [22] S. K. Lee, K. L. Sauer, S. J. Seltzer, O. Alem, and M. V. Romalis, Appl. Phys. Lett. **89**, 214106 (2006).
- [23] W. Wasilewski, K. Jensen, H. Krauter, J. J. Renema, and E. S. Polzik, Phys. Rev. Lett. **104**, 133601 (2010).